arXiv:quant-ph/0012036v2 11 Dec 2000

Covariant geometric quantization of non-relativistic Hamiltonian mechanics

GIOVANNI GIACHETTA$^{\dagger 1}$, LUIGI MANGIAROTTI$^{\dagger 2}$ AND GENNADI SARDANASHVILY$^{\ddagger 3}$

† Department of Mathematics and Physics, University of Camerino, 62032 Camerino (MC), Italy
‡ Department of Theoretical Physics, Physics Faculty, Moscow State University, 117234 Moscow, Russia

Abstract. We provide geometric quantization of the vertical cotangent bundle V^*Q equipped with the canonical Poisson structure. This is a momentum phase space of non-relativistic mechanics with the configuration bundle $Q \to \mathbf{R}$. The goal is the Schrödinger representation of V^*Q. We show that this quantization is equivalent to the fibrewise quantization of symplectic fibres of $V^*Q \to \mathbf{R}$ that makes the quantum algebra of non-relativistic mechanics an instantwise algebra. Quantization of the classical evolution equation defines a connection on this instantwise algebra, which provides quantum evolution in non-relativistic mechanics as a parallel transport along time.

1 Introduction

We study covariant geometric quantization of non-relativistic Hamiltonian mechanics subject to time-dependent transformations.

Its configuration space is a fibre bundle $Q \to \mathbf{R}$ equipped with bundle coordinates (t, q^k), $k = 1, \ldots, m$, where t is the Cartesian coordinate on the time axis $\mathbf{R}$ with affine transition functions $t' = t+$const. Different trivializations $Q \cong \mathbf{R} \times M$ of Q correspond to different non-relativistic reference frames. In contrary to all the existent quantizations of non-relativistic mechanics (e.g., [14, 19]), we do not fix a trivialization of Q.

The momentum phase space of non-relativistic mechanics is the vertical cotangent bundle V^*Q of $Q \to \mathbf{R}$, endowed with the holonomic coordinates (t, q^k, p_k). It is provided with the canonical Poisson structure

$$\{f, f'\}_V = \partial^k f \partial_k f' - \partial_k f \partial^k f', \qquad f, f' \in C^\infty(V^*Q), \tag{1}$$

---

[1] E-mail address: giachetta@campus.unicam.it

[2] E-mail address: mangiaro@camserv.unicam.it

[3] E-mail address: sard@grav.phys.msu.su

whose symplectic foliation coincides with the fibration $V^*Q \to \mathbf{R}$ [8, 12]. Given a trivialization

$$V^*Q \cong \mathbf{R} \times T^*M, \tag{2}$$

the Poisson manifold $(V^*Q, \{,\}_V)$ is isomorphic to the direct product of the Poisson manifold $\mathbf{R}$ with the zero Poisson structure and the symplectic manifold T^*M. An important peculiarity of the Poisson structure (1) is that the Poisson algebra $C^\infty(V^*Q)$ of smooth real functions on V^*Q is a Lie algebra over the ring $C^\infty(\mathbf{R})$ of functions of time alone.

Our goal is the geometric quantization of the Poisson bundle $V^*Q \to \mathbf{R}$, but it is not sufficient for quantization of non-relativistic mechanics.

The problem is that non-relativistic mechanics can not be described as a Poisson Hamiltonian system on the momentum phase space V^*Q. Indeed, a non-relativistic Hamiltonian $\mathcal{H}$ is not an element of the Poisson algebra $C^\infty(V^*Q)$. Its definition involves the cotangent bundle T^*Q of Q. Coordinated by $(q^0 = t, q^k, p_0 = p, p_k)$, the cotangent bundle T^*Q plays the role of the homogeneous momentum phase space of non-relativistic mechanics. It is equipped with the canonical Liouville form $\Xi = p_\lambda dq^\lambda$, the symplectic form $\Omega = d\Xi$, and the corresponding Poisson bracket

$$\{f, g\}_T = \partial^\lambda f \partial_\lambda f - \partial_\lambda f \partial^\lambda f', \qquad f, f' \in C^\infty(T^*Q).$$

Due to the one-dimensional canonical fibration

$$\zeta : T^*Q \to V^*Q, \tag{3}$$

the cotangent bundle T^*Q provides the symplectic realization of the Poisson manifold V^*Q, i.e.,

$$\zeta^*\{f, f'\}_V = \{\zeta^* f, \zeta^* f'\}_T$$

for all $f, f' \in C^\infty(V^*Q)$. A Hamiltonian on V^*Q is defined as a global section

$$h : V^*Q \to T^*Q, \qquad p \circ h = -\mathcal{H}(t, q^j, p_j), \tag{4}$$

of the one-dimensional affine bundle (3) [8, 12]. As a consequence (see Section 6), the evolution equation of non-relativistic mechanics is expressed into the Poisson bracket $\{,\}_T$ on T^*Q. It reads

$$\vartheta_{H^*}(\zeta^* f) = \{\mathcal{H}^*, \zeta^* f\}_T, \tag{5}$$

where ϑ_{H^*} is the Hamiltonian vector field of the function

$$\mathcal{H}^* = \partial_t \rfloor (\Xi - \zeta^* h^* \Xi)) = p + \mathcal{H} \tag{6}$$

on T^*Q.

Therefore, we need the compatible geometric quantizations both of the cotangent bundle T^*Q and the vertical cotangent bundle V^*Q such that the monomorphism

$$\zeta^* : (C^\infty(V^*Q), \{,\}_V) \to (C^\infty(T^*Q), \{,\}_T) \tag{7}$$

of the Poisson algebra on V^*Q to that on T^*Q is prolonged to a monomorphism of quantum algebras of V^*Q and T^*Q.

Recall that the geometric quantization procedure falls into three steps: prequantization, polarization and metaplectic correction (e.g., [3, 14, 19]). Given a symplectic manifold (Z, Ω) and the corresponding Poisson bracket $\{,\}$, prequantization associates to each element f of the Poisson algebra $C^\infty(Z)$ on Z a first order differential operator $\widehat{f}$ in the space of sections of a complex line bundle C over Z such that the Dirac condition

$$[\widehat{f}, \widehat{f'}] = -i\widehat{\{f, f'\}} \tag{8}$$

holds. Polarization of a symplectic manifold (Z, Ω) is defined as a maximal involutive distribution $\mathbf{T} \subset TZ$ such that $\mathrm{Orth}_\Omega \mathbf{T} = \mathbf{T}$, i.e.,

$$\Omega(\vartheta, \upsilon) = 0, \qquad \forall \vartheta, \upsilon \in \mathbf{T}_z, \qquad z \in Z. \tag{9}$$

Given the Lie algebra $\mathbf{T}(Z)$ of global sections of $\mathbf{T} \to Z$, let $\mathcal{A}_T \subset C^\infty(Z)$ denote the subalgebra of functions f whose Hamiltonian vector fields ϑ_f fulfill the condition

$$[\vartheta_f, \mathbf{T}(Z)] \subset \mathbf{T}(Z). \tag{10}$$

Elements of this subalgebra are only quantized. Metaplectic correction provides the pre-Hilbert space E_T where the quantum algebra $\mathcal{A}_T$ acts by symmetric operators. This is a certain subspace of sections of the tensor product $C \otimes \mathcal{D}_{1/2}$ of the prequantization line bundle $C \to Z$ and a bundle $\mathcal{D}_{1/2} \to Z$ of half-densities on Z. The geometric quantization procedure has been extended to Poisson manifolds [16, 17] and to Jacobi manifolds [7].

We show that standard prequantization of the cotangent bundle T^*Q (e.g., [3, 14, 19]) provides the compatible prequantization of the Poisson manifold V^*Q such that the monomorphism ζ^* (7) is prolonged to a monomorphism of prequantum algebras.

In contrast with the prequantization procedure, polarization of T^*Q need not imply a compatible polarization of V^*Q, unless it includes the vertical cotangent bundle $V_\zeta T^*Q$

of the fibre bundle ζ (3), i.e., spans over vectors ∂^0. The canonical real polarization of T^*Q, satisfying the condition

$$V_\zeta T^*Q \subset \mathbf{T}, \qquad (11)$$

is the vertical polarization. It coincides with the vertical tangent bundle VT^*Q of T^*Q, i.e., spans over all the vectors ∂^λ. We show that this polarization and the corresponding metaplectic correction of T^*Q induces the compatible quantization of the Poisson manifold V^*Q such that the monomorphism of Poisson algebras ζ^* (7) is prolonged to a monomorphism of quantum algebras of V^*Q and T^*Q. The quantum algebra $\mathcal{A}_V$ of V^*Q consists of functions on V^*Q which are at most affine in momenta p_k and have the Schrödinger representation in the space of half-densities on Q. It is essential that, since these operators does not contain the derivative with respect to time, the quantum algebra $\mathcal{A}_V$ is a $C^\infty(\mathbf{R})$-algebra.

We prove that the Schrödinger quantization of V^*Q yields geometric quantization of symplectic fibres of the Poisson bundle $V^*Q \to \mathbf{R}$ such that any quantum operator $\widehat{f}$ on V^*Q, restricted to the fibre V^*_tQ, $t \in \mathbf{R}$, coincides with the quantum operator on the symplectic manifold V^*_tQ of the function $f|_{V^*Q}$. Thus, the quantum algebra $\mathcal{A}_V$ of the Poisson bundle $V^*Q \to \mathbf{R}$ can be seen as the instantwise $C^\infty(\mathbf{R})$-algebra of its symplectic fibres. This agrees with the instantwise quantization of symplectic fibres $\{t\} \times T^*M$ of the direct product (2) in [14].

A fault of the Schrödinger representation on half-densities is that the Hamiltonian function $\mathcal{H}^*$ (6) does not belong to the quantum algebra $\mathcal{A}_T$ of T^*Q in general. A rather sophisticated solution of this problem for quadratic Hamiltonians has been suggested in [14]. However, the Laplace operator constructed in [14] does not fulfill the Dirac condition (8). If a Hamiltonian $\mathcal{H}$ is a polynomial of momenta p_k, one can represent it as an element of the universal enveloping algebra of the Lie algebra $\mathcal{A}_T$, but this representation is not necessarily globally defined.

In order to include a Hamiltonian function $\mathcal{H}^*$ (6) to the quantum algebra, one can choose the Hamiltonian polarization of T^*Q which contains the Hamiltonian vector field $\vartheta_{\mathcal{H}^*}$ of $\mathcal{H}^*$. However, it does not satisfy the condition (11) and does not define any polarization of the Poisson manifold V^*Q. This polarization is the necessary ingredient in a different variant of geometric quantization of V^*Q which is seen as a presymplectic manifold $(V^*Q, h^*\Omega)$. Given a trivialization (2), this quantization has been studied in [19]. In Section 5, its frame-covariant form is discussed.

Finally, since the quantum algebra $\mathcal{A}_V$ of the Poisson manifold V^*Q is a $C^\infty(\mathbf{R})$-algebra and since $\widehat{p} = -i\partial/\partial t$, quantization of the classical evolution equation (5) defines the connection

$$\nabla\widehat{f} = i[\widehat{\mathcal{H}}^*, \widehat{f}] \tag{12}$$

on the enveloping quantum algebra $\overline{\mathcal{A}}_V$ and describes quantum evolution in non-relativistic mechanics as a parallel transport along time.

2 Prequantization

Basing on the standard prequantization of the cotangent bundle T^*Q, we here construct the compatible prequantizations of the Poisson bundle $V^*Q \to \mathbf{R}$ and its symplectic leaves.

Recall the prequantization of T^*Q (e.g., [3, 14, 19]). Since its symplectic form Ω is exact and belongs to the zero de Rham cohomology class, the prequantization bundle is the trivial complex line bundle

$$C = T^*Q \times \mathbf{C} \to T^*Q, \tag{13}$$

whose Chern class c_1 is zero. Coordinated by $(q^\lambda, p_\lambda, c)$, it is provided with the admissible linear connection

$$A = dp_\lambda \otimes \partial^\lambda + dq^\lambda \otimes (\partial_\lambda + ip_\lambda c\partial_c) \tag{14}$$

with the strength form $F = -i\Omega$ and the Chern form

$$c_1 = \frac{i}{2\pi}F = \frac{1}{2\pi}\Omega.$$

The A-invariant Hermitian fibre metric on C is $g(c, c) = c\overline{c}$. The covariant derivative of sections s of the prequantization bundle C (13) relative to the connection A (14) along the vector field u on T^*Q takes the form

$$\nabla_u(s) = (u^\lambda\partial_\lambda - iu^\lambda p_\lambda)s. \tag{15}$$

Given a function $f \in C^\infty(T^*Q)$, the covariant derivative (15) along the Hamiltonian vector field

$$\vartheta_f = \partial^\lambda f\partial_\lambda - \partial_\lambda f\partial^\lambda, \qquad \vartheta_f\rfloor\Omega = -df$$

of f reads

$$\nabla_{\vartheta_f} = \partial^\lambda f(\partial_\lambda - ip_\lambda) - \partial_\lambda f\partial^\lambda.$$

Then, in order to satisfy the Dirac condition (8), one assigns to each element f of the Poisson algebra $C^\infty(T^*Q)$ the first order differential operator

$$\widehat{f}(s) = -i(\nabla_{\vartheta_f} + if)s = [-i\vartheta_f + (f - p_\lambda \partial^\lambda f)]s \tag{16}$$

on sections $s \in C(T^*Q)$ of the prequantization bundle C (13). For instance, the prequantum operators (16) for local functions $f = p_\lambda$, $f = q^k$, a global function $f = t$, and the constant function $f = 1$ read

$$\widehat{p}_\lambda = -i\partial_\lambda, \qquad \widehat{q}^\lambda = i\partial^\lambda + q^\lambda, \qquad \widehat{1} = 1.$$

For elements f of the Poisson subalgebra $C^\infty(V^*Q) \subset C^\infty(T^*Q)$, the Kostant–Souriau formula (16) takes the form

$$\widehat{f}(s) = [-i(\partial^k f \partial_k - \partial_\lambda f \partial^\lambda) + (f - p_k \partial^k f)]s. \tag{17}$$

Turn now to prequantization of the Poisson manifold $(V^*Q, \{,\}_V)$. The Poisson bivector w of the Poisson structure on V^*Q reads

$$w = \partial^k \wedge \partial_k = -[w, u]_{\rm SN}, \tag{18}$$

where $[,]_{\rm SN}$ is the Schouten–Nijenhuis bracket and $u = p_k \partial^k$ is the Liouville vector field on the vertical cotangent bundle $V^*Q \to Q$. The relation (18) shows that the Poisson bivector w is exact and, consequently, has the zero Lichnerowicz–Poisson cohomology class [8, 17]. Therefore, let us consider the trivial complex line bundle

$$C_V = V^*Q \times \mathbf{C} \to V^*Q \tag{19}$$

such that the zero Lichnerowicz–Poisson cohomology class of w is the image of the zero Chern class c_1 of C_V under the cohomology homomorphisms

$$H^*(V^*Q, \mathbf{Z}) \to H^*_{\rm deRh}(V^*Q) \to H^*_{\rm LP}(V^*Q).$$

Since the line bundles C (13) and C_V (19) are trivial, C can be seen as the pull-back $\zeta^* C_V$ of C_V, while C_V is isomorphic to the pull-back h^*C of C with respect to a section h (4) of the affine bundle (3). Since $C_V = h^*C$ and since the covariant derivative of the connection A (14) along the fibres of ζ (3) is trivial, let us consider the pull-back

$$h^*A = dp_k \otimes \partial^k + dq^k \otimes (\partial_k + ip_k c\partial_c) + dt \otimes (\partial_t - i\mathcal{H} c\partial_c) \tag{20}$$

of the connection A (14) onto $C_V \to V^*Q$ [9]. This connection defines the contravariant derivative

$$\nabla_\phi s_V = \nabla_{w^\sharp\phi} s_V \qquad (21)$$

of sections s_V of $C_V \to V^*Q$ along one-forms ϕ on V^*Q, which corresponds to a contravariant connection A_V on the line bundle $C_V \to V^*Q$ [17]. It is readily observed that this contravariant connection does not depend on the choice of a section h. By virtue of the relation (21), the curvature bivector of A_V equals to $-iw$ [18], i.e., A_V is an admissible connection for the canonical Poisson structure on V^*Q. Then the Kostant–Souriau formula

$$\widehat{f}_V(s_V) = (-i\nabla_{\vartheta_V f} + f)s_V = [-i(\partial^k f \partial_k - \partial_k f \partial^k) + (f - p_k\partial^k f)]s_V \qquad (22)$$

defines prequantization of the Poisson manifold V^*Q.

In particular, the prequantum operators of functions $f \in C^\infty(\mathbf{R})$ of time alone reduces simply to multiplication $\widehat{f}_V s_V = f s_V$ by these functions. Consequently, the prequantum algebra $\widehat{C}^\infty(V^*Q)$ inherits the structure of a $C^\infty(\mathbf{R})$-algebra.

It is immediately observed that the prequantum operator $\widehat{f}_V$ (22) coincides with the prequantum operator $\widehat{\zeta^* f}$ (17) restricted to the pull-back sections $s = \zeta^* s_V$ of the line bundle C. Thus, prequantization of the Poisson algebra $C^\infty(V^*Q)$ on the Poisson manifold $(V^*Q, \{,\})$ is equivalent to its prequantization as a subalgebra of the Poisson algebra $C^\infty(T^*Q)$ on the symplectic manifold T^*Q.

The above prequantization of the Poisson manifold V^*Q yields prequantization of its symplectic leaves as follows.

Since $w^\sharp\phi = \phi^k\partial_k - \phi_k\partial^k$ is a vertical vector field on $V^*Q \to \mathbf{R}$ for any one-form ϕ on V^*Q, the contravariant derivative (21) defines a connection along each fibre V_t^*Q, $t \in \mathbf{R}$, of the Poisson bundle $V^*Q \to \mathbf{R}$. This is the pull-back

$$A_t = i_t^* h^* A = dp_k \otimes \partial^k + dq^k \otimes (\partial_\lambda + ip_k c\partial_c)$$

of the connection h^*A (20) on the pull-back bundle $i_t^* C_V \to V_t^*Q$ with respect to the imbedding $i_t : V_t^*Q \to V^*Q$. It is readily observed that this connection is admissible for the symplectic structure

$$\Omega_t = dp_k \wedge dq^k$$

on V_t^*Q, and provides prequantization of the symplectic manifold (V_t^*Q, Ω_t). The corresponding prequantization formula is given by the expression (22) where functions f and sections s_V are restricted to V_t^*Q. Thus, the prequantization (22) of the Poisson manifold V^*Q is a leafwise prequantization [18].

3 Polarization

Given compatible prequantizations of the cotangent bundle T^*Q, the Poisson bundle $V^*Q\to\mathbf{R}$ and its simplectic fibres, let us now construct their compatible polarizations.

Recall that, given a polarization $\mathbf{T}$ of a prequantum symplectic manifold (Z,Ω), the subalgebra $\mathcal{A}_T\subset C^\infty(Z)$ of functions f obeying the condition (10) is only quantized. Moreover, after further metaplectic correction, one consider a representation of this algebra in a quantum space E_T such that

$$\nabla_u e=0, \qquad \forall u\in\mathbf{T}(Z), \qquad e\in E_T. \tag{23}$$

Recall that by a polarization of a Poisson manifold $(Z,\{,\})$ is meant a sheaf $\mathbf{T}^*$ of germs of complex functions on Z whose stalks $\mathbf{T}^*_z$, $z\in Z$, are Abelian algebras with respect to the Poisson bracket $\{,\}$ [18]. One can also require that the algebras $\mathbf{T}^*_z$ are maximal, but this condition need not hold under pull-back and push-forward operations. Let $\mathbf{T}^*(Z)$ be the structure algebra of global sections of the sheaf $\mathbf{T}^*$; it is also called a Poisson polarization [16, 17]. A quantum algebra $\mathcal{A}_T$ associated to the Poisson polarization $\mathbf{T}^*$ is defined as a subalgebra of the Poisson algebra $C^\infty(Z)$ which consists of functions f such that

$$\{f,\mathbf{T}^*(Z)\}\subset\mathbf{T}^*(Z).$$

Polarization of a symplectic manifold yields its maximal Poisson polarization, and *vice versa*.

There are different polarizations of the cotangent bundle T^*Q. We will consider those polarizations of T^*Q whose direct image as Poisson polarizations onto V^*Q with respect to the morphism ζ (3) are polarizations of the Poisson manifold V^*Q. This takes place if the germs of polarization $\mathbf{T}^*$ of the Poisson manifold $(T^*Q,\{,\}_T)$ are constant along the fibres of the fibration ζ (3) [18], i.e., are germs of functions independent of the momentum coordinate $p_0=p$. It means that the corresponding polarization $\mathbf{T}$ of the symplectic manifold T^*Q is vertical with respect to the fibration $T^*Q\to\mathbf{R}$, i.e., obeys the condition (11). A short calculation shows that, in this case, the associated quantum algebra $\mathcal{A}_T$ consists of functions $f\in C^\infty(T^*Q)$ which are at most affine in the momentum coordinate p_0. Moreover, given such a polarization, the equality (23) implies the equality

$$\nabla_{u_0\partial^0}e=0, \qquad e\in E_T,$$

for any vertical vector field $u_0\partial^0$ on the fibre bundle $T^*Q\to V^*Q$. Then the prequantization formulas (17) and (22) for the Poisson algebra $C^\infty(V^*Q)$ coincide on quantum spaces, i.e., the monomorphism ζ^* (7) is prolonged to monomorphism of quantum algebras of V^*Q and T^*Q.

The vertical polarization VT^*Q of T^*Q obeys the condition (11). It is a strongly admissible polarization, and its integral manifolds are fibres of the cotangent bundle $T^*Q \to Q$. One can verify easily that the associated quantum algebra $\mathcal{A}_T$ consists of functions on T^*Q which are at most affine in momenta p_λ. The quantum space E_T associated to the vertical polarization obeys the condition

$$\nabla_{u^\lambda\partial_\lambda} e = 0, \qquad \forall e \in E_T. \tag{24}$$

Therefore, the operators of the quantum algebra $\mathcal{A}_T$ on this quantum space read

$$f = a^\lambda(q^\mu)p_\lambda + b(q^\mu), \qquad \widehat{f} = -i\nabla_{a^\lambda\partial_\lambda} + b. \tag{25}$$

This is the Schrödinger representation of T^*Q.

The vertical polarization of T^*Q defines the maximal polarization $\mathbf{T}^*$ of the Poisson manifold V^*Q which consists of germs of functions constant on the fibres of $V^*Q \to Q$. The associated quantum space E_V obeys the condition

$$\nabla_{u^k\partial_k} e = 0, \qquad \forall e \in E_V. \tag{26}$$

The quantum algebra $\mathcal{A}_V$ corresponding to this polarization of V^*Q consists of functions on V^*Q which are at most affine in momenta p_k. Their quantum operators read

$$f = a^\lambda(q^\mu)p_k + b(q^\mu), \qquad \widehat{f} = -i\nabla_{a^k\partial_k} + b. \tag{27}$$

This is the Schrödinger representation of V^*Q.

In turn, each symplectic fibre V_t^*Q, $t \in \mathbf{R}$, of the Poisson bundle $V^*Q \to \mathbf{R}$ is provided with the pull-back polarization $\mathbf{T}_t^* = i_t^*\mathbf{T}^*$ with respect to the Poisson morphism $i_t : V_t^*Q \to V^*Q$. The corresponding distribution $\mathbf{T}_t$ coincides with the vertical tangent bundle of the fibre bundle $V_t^*Q \to Q_t$. The associated quantum algebra $\mathcal{A}_t$ consists of functions on V^*Q_t which are at most affine in momenta p_k, while the quantum space E_t obeys the condition similar to (26). Therefore, the representation of the quantum algebra $\mathcal{A}_t$ takes the form (27). It follows that the Schrödinger representation the Poisson bundle $V^*Q \to \mathbf{R}$ is a fibrewise representation.

4 Metaplectic correction

To complete the geometric quantization procedure of V^*Q, let us consider the metaplectic correction of the Schrödinger representations of T^*Q and V^*Q.

The representation of the quantum algebra $\mathcal{A}_T$ (25) can be defined in the subspace of sections of the line bundle $C \to T^*Q$ which fulfill the relation (24). This representation reads

$$f = a^\lambda(q^\mu)p_\lambda + b(q^\mu), \qquad \widehat{f} = -ia^\lambda\partial_\lambda + b. \tag{28}$$

Therefore, it can be restricted to the sections s of the pull-back line bundle $C_Q = \widehat{0}^*C \to Q$ where $\widehat{0}$ is the canonical zero section of the cotangent bundle $T^*Q \to Q$. However, this is not yet a representation in a Hilbert space.

Let Q be an oriented manifold. Applying the general metaplectic technique [3, 19], we come to the vector bundle $\mathcal{D}_{1/2} \to Q$ of complex half-densities on Q with the transition functions $\rho' = J^{-1/2}\rho$, where J is the Jacobian of the coordinate transition functions on Q. Since $C_Q \to Q$ is a trivial bundle, the tensor product $C_Q \otimes \mathcal{D}_{1/2}$ is isomorphic to $\mathcal{D}_{1/2}$. Therefore, the quantization formula (28) can be extended to sections of the half-density bundle $\mathcal{D}_{1/2} \to Q$ as

$$f = a^\lambda(q^\mu)p_\lambda + b(q^\mu), \qquad \widehat{f}\rho = (-i\mathbf{L}_{a^\lambda\partial_\lambda} + b)\rho = (-ia^\lambda\partial_\lambda - \frac{i}{2}\partial_\lambda(a^\lambda) + b)\rho, \tag{29}$$

where $\mathbf{L}$ denotes the Lie derivative. The second term in the right-hand side of this formula is a metaplectic correction. It makes the operator $\widehat{f}$ (29) symmetric with respect to the Hermitian form

$$\langle\rho_1|\rho_2\rangle = \left(\frac{1}{2\pi}\right)^m \int_Q \rho_1\overline{\rho}_2$$

on the pre-Hilbert space E_T of sections of $\mathcal{D}_{1/2}$ with compact support. The completion $\overline{E}_T$ of E_T provides a Hilbert space of the Schrödinger representation of the quantum algebra $\mathcal{A}_T$, where the operators (29) are essentially self-adjoint, but not necessarily bounded. Of course, functions of compact support on the time axis $\mathbf{R}$ have a limited physical application, but we can always restrict our consideration to some bounded interval of $\mathbf{R}$.

Since, in the case of the vertical polarization, there is a monomorphism of the quantum algebra $\mathcal{A}_V$ to the quantum algebra $\mathcal{A}_T$, one can define the Schrödinger representation of $\mathcal{A}_V$ by the operators

$$f = a^k(q^\mu)p_k + b(q^\mu), \qquad \widehat{f}\rho = (-ia^k\partial_k - \frac{i}{2}\partial_k(a^k) + b)\rho \tag{30}$$

in the same space of complex half-densities on Q as that of $\mathcal{A}_T$. Moreover, this representation preserves the structure of $\mathcal{A}_V$ as a $C^\infty(\mathbf{R})$-algebra.

The metaplectic correction of the Poisson bundle $V^*Q \to \mathbf{R}$ also provides the metaplectic corrections of its symplectic leaves as follows. It is readily observed that the Jacobian J restricted to a fibre Q_t, $t \in \mathbf{R}$, is a Jacobian of coordinate transformations on Q_t. Therefore, any half-density ρ on Q, restricted to Q_t, is a half-density on Q_t. Then the representation (30) restricted to a fibre V^*_tQ is exactly the metaplectic correction of the Schrödinger representation of the symplectic manifold V^*_tQ on half-densities on Q_t.

Thus, the quantum algebra $\mathcal{A}_V$ of the Poisson bundle $V^*Q \to \mathbf{R}$, given by the operators (30), can be seen as the instantwise algebra of operators on its symplectic fibres.

The representation (30) can be extended locally to functions on T^*Q which are polynomials of momenta p_λ. These functions can be represented by elements of the universal enveloping algebra $\overline{\mathcal{A}}_T$ of the Lie algebra $\mathcal{A}_T$, but this representation is not necessarily globally defined. For instance, a generic quadratic Hamiltonian

$$\mathcal{H} = a^{jk}(q^\lambda)p_jp_k + b^k(q^\lambda)p_k + c(q^\lambda) \tag{31}$$

leads to the Hamiltonian function $\mathcal{H}^* = p + \mathcal{H}$ which is not an element of $\overline{\mathcal{A}}_T$ because of the quadratic term. This term can be quantized only locally, unless the Jacobian of the coordinate transition functions on Q is independent of fibre coordinates q^k on Q.

5 Presymplectic quantization

As was mentioned above, to include a Hamiltonian function $\mathcal{H}^*$ (6) to the quantum algebra, one can choose the Hamiltonian polarization of T^*Q. This polarization accompanies a different approach to geometric quantization of the momentum phase space of non-relativistic mechanics V^*Q which is considered as a presymplectic manifold. In comparison with the above one, this quantization does not lead to quantization of the Poisson algebra of functions on V^*Q as follows.

Every global section h (4) of the affine bundle ζ (3) yields the pull-back Hamiltonian form

$$H = h^*\Xi = p_kdq^k - \mathcal{H}dt \tag{32}$$

on V^*Q. With respect to a trivialization (2), the form H is the well-known integral invariant of Poincaré–Cartan [1]. Given a Hamiltonian form H (32), there exists a unique Hamiltonian connection

$$\gamma_H = \partial_t + \partial^k\mathcal{H}\partial_k - \partial_k\mathcal{H}\partial^k \tag{33}$$

on the fibre bundle $V^*Q \to \mathbf{R}$ such that

$$\gamma_H \rfloor dH = 0 \tag{34}$$

[8, 12]. It defines the Hamilton equations on V^*Q.

A glance at the equation (34) shows that one can think of the Hamiltonian connection γ_H as being the Hamiltonian vector field of a zero Hamiltonian with respect to the presymplectic form dH on V^*Q. Therefore, one can study geometric quantization of the presymplectic manifold (V^*Q, dH).

Usually, geometric quantization is not applied directly to a presymplectic manifold (Z, ω), but to a symplectic manifold (Z', ω') such that the presymplectic form ω is a pull-back of the symplectic form ω'. Such a symplectic manifold always exists. The following two possibilities are usually considered: (i) (Z', ω') is a reduction of (Z, ω) along the leaves of the characteristic distribution of the presymplectic form ω of constant rank [6, 15], and (ii) there is a coisotropic imbedding of (Z, ω) to (Z', ω') [4, 5].

In application to (V^*Q, dH), the reduction procedure however meets difficulties. Since the kernel of dH is generated by the vectors $(\partial_t,\ \partial_k\mathcal{H}\partial^k - \partial^k\mathcal{H}\partial_k,\ k = 1, \ldots, m)$, the presymplectic form dH in physical models is almost never of constant rank. Therefore, one has to provide an exclusive analysis of each physical model, and to cut out a certain subset of V^*Q in order to use the reduction procedure.

The second variant of geometric quantization of the presymplectic manifold (V^*Q, dH) seems more attractive because the section h (4) is a coisotropic imbedding. Indeed, the tangent bundle TN_h of the closed imbedded submanifold $N_h = h(V^*Q)$ of T^*Q consists of the vectors

$$u = -(u^\mu \partial_\mu \mathcal{H} + u_j \partial^j \mathcal{H})\partial^0 + u_k \partial^k + u^\mu \partial_\mu. \tag{35}$$

Let us prove that the orthogonal distribution $\mathrm{Orth}_\Omega TN_h$ of TN_h with respect to the symplectic form Ω belongs to TN_h. By definition, it consists of the vectors $u \in T_zT^*Q$, $z \in T^*Q$, such that

$$u \rfloor v \rfloor \Omega = 0, \qquad \forall v \in T_zN_h.$$

A simple calculation shows that these vectors obey the conditions

$$-u^0 \partial^i \mathcal{H} + u^i = 0, \qquad u^0 \partial_\mu - u_\mu = 0$$

and, consequently, take the form (35).

The image $N_h = h(V^*Q)$ of the coisotropic imbedding h is given by the constraint

$$\mathcal{H}^* = p + \mathcal{H}(t, q^k, p_k) = 0.$$

Then the geometric quantization of the presymplectic manifold (V^*Q, dH) consists in geometric quantization of the cotangent bundle T^*Q and setting the quantum constraint condition

$$\widehat{\mathcal{H}^*}\psi = 0$$

on physically admissible quantum states. This condition implies that $\widehat{\mathcal{H}^*}$ belongs to the quantum algebra of T^*Q. It takes place if the above mentioned Hamiltonian polarization of T^*Q, which contains the Hamiltonian vector field

$$\vartheta_{\mathcal{H}^*} = \partial_0 + \partial^k\mathcal{H}\partial_k - \partial_k\mathcal{H}\partial^k, \tag{36}$$

is used.

Such a polarization of T^*Q always exists. Indeed, any section h (4) of the affine bundle $T^*Q \to V^*Q$ defines the splitting

$$a_\lambda\partial^\lambda = a_k(\partial^k - \partial^k\mathcal{H}\partial^0) + (a_0 + a_k\partial^k\mathcal{H})\partial^0$$

of the vertical tangent bundle VT^*Q of $T^*Q \to Q$. One can justify this fact, e.g., by inspection of the coordinate transformation law. Then elements $(\partial^k - \partial^k\mathcal{H}\partial^0$, $k = 1, \ldots, m)$, and the values of the Hamiltonian vector field $\vartheta_{\mathcal{H}^*}$ (36) obey the polarization condition (9) and generates a polarization of T^*Q. It is clear that the Hamiltonian polarization does not satisfy the condition (11), and does not define any polarization of the Poisson manifold V^*Q.

Nevertheless, given a trivialization (2), symplectic fibres V_t^*Q, $t \in \mathbf{R}$, of the Poisson bundle $V^*Q \to \mathbf{R}$ can be provided with the Hamiltonian polarization $\mathbf{T}_t$ generated by vectors $\partial_k\mathcal{H}\partial^k - \partial^k\mathcal{H}\partial_k$, $k = 1, \ldots, m$, except the points where

$$d\mathcal{H} = \partial_k\mathcal{H}dq^k + \partial^k\mathcal{H}dp_k = 0.$$

This is a standard polarization in conservative Hamiltonian mechanics of one-dimensional systems, but it requires an exclusive analysis of each model.

6 Classical and quantum evolution equations

Turn now to the evolution equation in classical and quantum non-relativistic mechanics.

Given a Hamiltonian connection γ_H on the momentum phase space V^*Q, let us consider the Lie derivative

$$\mathbf{L}_{\gamma_H} f = \gamma_H \rfloor df = (\partial_t + \partial^k\mathcal{H}\partial_k - \partial_k\mathcal{H}\partial^k)f \tag{37}$$

of a function $f \in C^\infty(V^*Q)$ along γ_H. This equality is the evolution equation in classical non-relativistic Hamiltonian mechanics. Substituting a solution of the Hamilton equations in its right-hand side, one obtains the time evolution of f along this solution. Given a trivialization (2), the evolution equation (37) can be written as

$$\mathbf{L}_\gamma f = \partial_t f + \{\mathcal{H}, f\}_V.$$

However, taken separately, the terms in its right-hand side are ill-behaved under time-dependent transformations. Let us bring the evolution equation into the frame-covariant form.

The affine bundle ζ (3) is modelled over the trivial line bundle $V^*Q \times \mathbf{R} \to V^*Q$. Therefore, Hamiltonian forms H constitute an affine space modelled over the vector space $C^\infty(V^*Q)$. To choose a centre of this affine space, let us consider a connection

$$\Gamma = \partial_t + \Gamma^k\partial_k \tag{38}$$

on the configuration bundle $Q \to \mathbf{R}$. By definition, it is a section of the affine bundle (3), and yields the Hamiltonian form

$$H_\Gamma = \Gamma^*\Xi = p_k dq^k - \mathcal{H}_\Gamma dt, \qquad \mathcal{H}_\Gamma = p_k\Gamma^k dt.$$

Its Hamiltonian connection is the canonical lift

$$V^*\Gamma = \partial_t + \Gamma^i\partial_i - p_i\partial_j\Gamma^i\partial^j$$

of the connection Γ onto $V^*Q \to \mathbf{R}$. Then any Hamiltonian form H (32) on the momentum phase space V^*Q admits the splittings

$$H = H_\Gamma - \widetilde{\mathcal{H}}_\Gamma dt, \qquad \widetilde{\mathcal{H}}_\Gamma = \mathcal{H} - p_k\Gamma^k, \tag{39}$$

where $\widetilde{\mathcal{H}}_\Gamma$ is a function on V^*Q. The physical meaning of this splitting becomes clear due to the fact that every trivialization of $Q \to \mathbf{R}$ yields a complete connection Γ on Q, and *vice versa* [8, 9]. From the physical viewpoint, the vertical part of this connection Γ

(38) can be seen as a velocity of an "observer", and Γ characterizes a reference frame in non-relativistic time-dependent mechanics [8, 11, 12]. Then one can show that $\widetilde{\mathcal{H}}_\Gamma$ in the splitting (39) is the energy function with respect to this reference frame [2, 8, 12].

Given the splitting (39), the evolution equation can be written in the frame-covariant form

$$\mathbf{L}_{\gamma_H} f = V^*\Gamma\rfloor H + \{\widetilde{\mathcal{H}}_\Gamma, f\}_V.$$

However, the first term in its right-hand side is not reduced to the Poisson bracket on V^*Q, and is not quantized in the framework of geometric quantization of the Poisson manifold V^*Q. To bring the right-hand side of the evolution equation into a Poisson bracket alone, let us consider the pull-back ζ^*H of the Hamiltonian form $H = h^*\Xi$ onto the cotangent bundle T^*Q. It is readily observed that the difference $\Xi - \zeta^*H$ is a horizontal 1-form on $T^*Q \to \mathbf{R}$, and we obtain the function (6) on T^*Q. Then the relation

$$\zeta^*(\mathbf{L}_{\gamma_H} f) = \vartheta_{H^*}(\zeta^* f) = \{\mathcal{H}^*, \zeta^* f\}_T,$$

holds for any element f of the Poisson algebra $C^\infty(V^*Q)$ [10].

Since the quantum algebra $\mathcal{A}_V$ of the Poisson manifold V^*Q can be seen as the instantwise algebra, one can quantize the evolution equation (5) as follows.

Given the Hamiltonian function $\mathcal{H}^*$ (6), let $\widehat{\mathcal{H}}^*$ be the corresponding quantum operator written, e.g., as an element of universal enveloping algebra of the quantum algebra $\mathcal{A}_T$. The bracket (12) defines a derivation of the enveloping algebra $\overline{\mathcal{A}}_V$ of the quantum algebra $\mathcal{A}_V$, which is also a $C^\infty(\mathbf{R})$-algebra. Moreover, since $\widehat{p} = -i\partial/\partial t$, the derivation (12) obeys the Leibniz rule

$$\nabla(r(t)\widehat{f}) = \partial_t r(t)\widehat{f} + r(t)\nabla\widehat{f}.$$

Therefore, it is a connection on the $C^\infty(\mathbf{R})$-algebra $\overline{\mathcal{A}}_V$, and defines quantum evolution of $\overline{\mathcal{A}}_V$ as a parallel transport along time [9, 13].

References

[1] V.Arnold, *Mathematical Methods of Classical Mechanics* (Springer, Berlin, 1978).

[2] A.Echeverría Enríquez, M.Muñoz Lecanda and N.Román Roy, Non-standard connections in classical mechanics, *J. Phys. A* **28** (1995) 5553.

[3] A.Echeverría Enríquez, M.Muñoz Lecanda, N.Román Roy, and C.Victoria-Monge, Mathematical foundations of geometric quantization, *Extracta Math.* **13** (1998) 135.

[4] M.Gotay and J.Śniatycki, On the quantization of presymplectic dynamical systems via coisotropic imbeddings, *Commun. Math. Phys.* **82** (1981) 377.

[5] M.Gotay, On coisotropic imbeddings of presymplectic manifolds, *Proc. Amer. Math. Soc.* **84** (1982) 111.

[6] M.Gotay, Constraints, reduction and quantization, *J. Math. Phys.* (1986) 2051.

[7] M. de León, J.Marrero and E.Padron, On the geometric quantization of Jacobi manifolds, *J. Math. Phys.* **38** (1997) 6185.

[8] L.Mangiarotti and G.Sardanashvily, *Gauge Mechanics* (World Scientific, Singapore, 1998).

[9] L.Mangiarotti and G.Sardanashvily, *Connections in Classical and Quantum Field Theory* (World Scientific, Singapore, 2000).

[10] L.Mangiarotti and G.Sardanashvily, Constraints in Hamiltonian time-dependent mechanics, *J. Math. Phys.* **41** (2000) 2858

[11] E.Massa and E.Pagani, Jet bundle geometry, dynamical connections and the inverse problem of Lagrangian mechanics, *Ann. Inst. Henri Poincaré* **61** (1994) 17.

[12] G.Sardanashvily, Hamiltonian time-dependent mechanics, *J. Math. Phys.* **39** (1998) 2714.

[13] G.Sardanashvily, Classical and quantum mechanics with time-dependent parameters, *J. Math. Phys.* **41** (2000) 5245.

[14] J.Śniatycki, *Geometric Quantization and Quantum Mechanics* (Springer-Verlag, Berlin, 1980).

[15] I.Vaisman, Geometric quantization on presymplectic manifold, *Monatsh. Math.* **96** (1983) 293.

[16] I.Vaisman, On the geometric quantization of Poisson manifolds, *J. Math. Phys.* **32** (1991) 3339.

[17] I.Vaisman, *Lectures on the Geometry of Poisson Manifolds* (Birkhäuser Verlag, Basel, 1994).

[18] I.Vaisman, On the geometric quantization of the symplectic leaves of Poisson manifolds, *Diff. Geom. Appl.* **7** (1997) 265.

[19] N.Woodhouse, *Geometric Quantization* (Clarendon Press, Oxford, 1980) (2nd ed. 1992).